

Article

Validation of the simplified modified Rankin scale for stroke trials: Experience from the ENCHANTED alteplase-dose arm

Chen, Xiaoying, Li, Jingwei, Anderson, Craig S, Lindley, Richard I, Hackett, Maree, Robinson, Thompson, Lavados, Pablo M, Wang, Xia, Arima, Hisatomi, Chalmers, John and Delcourt, Candice

Available at <http://clock.uclan.ac.uk/31527/>


Chen, Xiaoying, Li, Jingwei, Anderson, Craig S, Lindley, Richard I, Hackett, Maree ORCID: 0000-0003-1211-9087, Robinson, Thompson, Lavados, Pablo M, Wang, Xia, Arima, Hisatomi et al (2021) Validation of the simplified modified Rankin scale for stroke trials: Experience from the ENCHANTED alteplase-dose arm. International Journal of Stroke, 16 (2). pp. 222-228. ISSN 1747-4930

It is advisable to refer to the publisher's version if you intend to cite from the work.
<http://dx.doi.org/10.1177/1747493019897858>

For more information about UCLan's research in this area go to
<http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to
<http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.



Validation of the simplified modified Rankin scale for stroke trials: experience from the ENCHANTED alteplase-dose arm

Journal:	<i>International Journal of Stroke</i>
Manuscript ID	IJS-07-19-7266.R2
Manuscript Type:	Research
Date Submitted by the Author:	24-Oct-2019
Complete List of Authors:	Chen, Xiaoying; The George Institute for Global Health,

	<p>li, jINGWEI; George Institute for Global Health, Anderson, Craig; University of New South Wales, The George Institute for Global Health, Faculty of Medicine; Peking University Health Sciences Centre, ; Royal Prince Alfred Hospital</p> <p>Lindley, Richard; The University of Sydney Westmead Clinical School</p> <p>Hackett, Maree; The George Institute for Global Health, Neurological & Mental Health Division, The University of Sydney; The University of Central Lancashire, School of Health</p> <p>Robinson, Tom</p> <p>Lavados, Pablo; Universidad del Desarrollo, Departamento de Neurologia y Psiquiatria</p> <p>Wang, Xia; The George Institute for Global Health , Neurological & Mental Health Division</p> <p>Arima, Hisatomia; The George Institute for Global Health, NMH; Fukuoka University, Department of Preventive Medicine and Public Health, Faculty of Medicine</p> <p>Chalmers, John</p> <p>Delcourt, Candice; The George Institute,</p>
Keywords:	simplified modified Rankin scale questionnaire, modified Rankin scale, ischemic stroke, Clinical trial, health outcome, Thrombolysis

Validation of the simplified modified Rankin scale for stroke trials: experience from the ENCHANTED alteplase-dose arm

Xiaoying Chen BPharm BMgt,^{1,2} Jingwei Li MD PhD,^{1,3,4} Craig S. Anderson MD PhD,^{1,5,6} Richard I. Lindley MD,⁷ Maree L Hackett PhD,^{1,2,8} Thompson Robinson MD,^{9,10} Pablo M. Lavados MD MPH,^{11,12} Xia Wang PhD,¹ Hisatomi Arima PhD,^{1,13} John Chalmers MD PhD,¹ Candice Delcourt MD PhD,^{1,2,6} for the ENCHANTED Investigators*

¹The George Institute for Global Health, Faculty of Medicine, University of New South Wales, Sydney, Australia.

²Faculty of Medicine and Health, University of Sydney, NSW, Australia

³Department of Cardiology, People's Liberation Army General Hospital, Beijing, China

⁴Department of Cardiology, Xinqiao Hospital, Third Military Medical University, Chongqing, China

⁵The George Institute China at Peking University Health Science Center, Beijing, PR China

⁶Neurology Department, Royal Prince Alfred Hospital, Sydney Health Partners, Sydney, NSW, Australia

⁷Westmead Clinical School, University of Sydney, NSW, Australia;

⁸Faculty of Health and Wellbeing, University of Central Lancashire, UK

⁹Department of Cardiovascular Sciences, University of Leicester, Leicester, UK

¹⁰NIHR Biomedical Research Centre, Leicester, UK

¹¹Departamento de Ciencias Neurológicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile

¹²Unidad de Neurología Vascular, Servicio de Neurología, Departamento de Neurología y Psiquiatría, Clínica Alemana de Santiago, Facultad de Medicina, Universidad del Desarrollo, Santiago, Chile

¹³Department of Preventive Medicine and Public Health, Faculty of Medicine, Fukuoka University, Fukuoka, Japan

Correspondence to:

Professor Craig Anderson,

The George Institute for Global Health, University of New South Wales

PO Box M201, Missenden Road, NSW 2050, Sydney, Australia

E: canderson@georgeinstitute.org.au; T: +61 2 9993 4521 ; F : +61 2 9993 4502

Word Count: abstract 265; body 2398

Key words: simplified modified Rankin scale questionnaire, modified Rankin scale, ischemic stroke, clinical trial, health outcome

Cover title: Validity of simplified mRS questionnaire

Supp. Tables: 3

Figures: 2

Subject Terms: quality and outcomes, ischemic stroke
List of tables and figures:

- [Supp.](#) Table 1. Correlation between smRSq and mRS scores at Day 90
- [Supp.](#) Table 2. Independent predictors of smRSq and mRS at Day 90
- [Supp.](#) Table 3. Comparison of treatment effects using mRS and smRSq in the alteplase-dose arm of the ENCHANTED trial
- Figure 1. Bubble plot of agreement between smRSq and mRS at Day 90
- Figure 2. ROC curves for predictive models of mRS and smRSq at Day 90.

For Review Only

Abstract

Background and Aims: The structured, simplified modified Rankin scale questionnaire (smRSq) may increase reliability over the interrogative approach to scoring the modified Rankin scale (mRS) in acute stroke research and practice. During the conduct of the alteplase-dose-arm of the international ENhanced Control of Hypertension ANd Thrombolysis Stroke stuDy (ENCHANTED), we had an opportunity to compare each of these approaches to outcome measurement.

Methods: Baseline demographic data were recorded together with the National Institutes of Health Stroke Scale (NIHSS). Follow-up measures obtained at 90 days included mRS, smRSq, and the 5-Dimension European Quality of life scale (EQ-5D). Agreements between smRSq and mRS were assessed with the Kappa statistic. Multiple logistic regression was used to identify baseline predictors of Day 90 smRSq and mRS scores. Treatment effects, based on Day 90 smRSq/mRS scores were tested in logistic and ordinal logistic regression models.

Results: SmRSq and mRS scores had good agreement (weighted Kappa 0.79, 95% confidence interval [CI] 0.78-0.81), whilst variables of age, atrial fibrillation, diabetes mellitus, pre-morbid mRS (1 vs. 0), baseline NIHSS scores and imaging signs of cerebral ischemia, similarly predicted their scores. Odds ratios for death or disability, and ordinal shift, 90 day mRS scores using smRSq were 1.05 (95% CI 0.91-1.20; one-sided $p=0.23$ for noninferiority) and 0.98 (95% CI 0.87-1.11; $P=0.02$ for noninferiority), similar to those using mRS.

Conclusions: This study demonstrates the utility of the smRSq in a large, ethnically diverse clinical trial population. Scoring of the smRSq shows adequate agreement with the standard mRS, thus confirming it is a reliable, valid and useful alternative measure of functional status after acute ischemic stroke.

Clinical Trial registration URL: <http://www.clinicaltrials.gov>. Unique identifier: NCT01422616

Introduction and Aims

The modified Rankin scale (mRS) is the most popular assessment tool for measuring overall functional status in patients who have suffered a stroke or other form of neurological disability,¹ both in clinical practice and research.^{2, 3} However, due to criticism being raised over subjectivity in aspects of its categorization/scoring,⁴ Bruno et al. developed the short, structured, simplified modified Rankin scale questionnaire (smRSq)^{5, 6} which has been shown to correlate with the size of the ischemic lesion,⁶ health-related quality of life,⁷ and neurological severity⁸ in small single center studies. The smRSq has also shown good reliability and validity in Chinese stroke patients.⁹ However, it has not been validated in a broader population or in the context of international research where the mRS remains the gold standard method of outcome assessment. We aimed to compare scores on the mRS and smRSq, their predictor variables, their correlation with neurological impairment on the National Institutes of Health Stroke Scale (NIHSS) and health-related quality of life on the 5-Dimension European Quality of life scale (EQ-5D), and treatment effects using them as outcome measures, among participants of the alteplase-dose arm of the Enhanced Control of Hypertension and Thrombolysis Stroke study (ENCHANTED).

Methods

Study design

ENCHANTED was an international, multicenter, quasi-factorial, prospective, randomized, open, blinded outcome assessed, clinical trial that assessed the effectiveness of low versus standard dose intravenous alteplase, and intensive versus guideline-recommended blood pressure (BP) management, in thrombolysis-eligible patients with acute ischemic stroke, the details of which are described elsewhere.^{10, 11} In brief, the first arm of the trial assessed 0.6 mg/kg compared to 0.9 mg/kg alteplase in 3310 patients (age ≥ 18 years) within 4.5 hours of

the onset of symptoms and followed up these patients to 90 days. The primary endpoint was death or disability defined by scores of 2 to 6 on the mRS. The trial was approved by local ethics committees and regulatory bodies, and written informed consent was obtained from the patient or an appropriate surrogate. The trial is registered at ClinicalTrials.gov (NCT01422616).

Measures

Demographics, clinical characteristics including the severity of neurological impairment on the NIHSS, were recorded in participants at the time of enrolment (baseline). The trial excluded patients with pre-morbid functional impairment (mRS scores >1) but collected estimated pre-morbid mRS (0 or 1) for those included. Signs of cerebral ischemia on brain imaging, and any evidence of proximal vessel occlusion on computed tomographic angiography (CTA) or magnetic resonance angiography (MRA), were reported by clinicians. Assessors with a health professional background (doctors, nurses or scientists) blind to treatment allocation and who had received in-person and online training (<https://secure.trainingcampus.net>), recorded mRS and smRSq scores by telephone or face-to-face interview in patients or a suitable proxy at 28 and 90 days post-randomisation. These outcome assessors had no mandatory training in the use of smRSq. They were advised to first assess patients with the mRS and then immediately administer the smRSq, as listed on the case report form. The 7-item mRS covers no symptoms (score 0), symptoms but no significant disability (1), slight disability (2), moderate disability (3), moderately severe disability (4), severe disability (5), and death (6). The smRSq takes on average 1.7 minutes to administer,⁷ and represents mRS items through yes/no answers to 5 questions addressing key functional states: living alone without any help from another person for bathing, toileting, shopping, preparing or getting meals, and managing finances; doing everything as before the stroke; being back to pre-stroke status; walking without help from another person; and being bedridden or needing constant supervision. The EQ-5D, which was

also administered directly in a patient or proxy at 28 and 90 days, defines the state of general health across five dimensions (mobility, self-care, usual activities, pain/discomfort, and anxiety/depression) with three levels of responses within each dimension (no problems, some/moderate problems, and severe problems). The EQ-5D utility score integrates the ratings of the 5 dimensions into a single score, calculated using population-based preference weights for each subscale. The weights used in the present analyses were derived from a study based on a representative sample of the UK population.¹² Utility scores express HRQoL quantitatively as a fraction of perfect health, with a score of 1 representing perfect health, a score of 0 representing death, and negative scores (minimum score -0.594) representing health states considered worse than death.¹³

Statistical analysis

Strength of agreement on ordinal analysis¹⁴ of the smRSq and mRS at Day 90 were assessed through Cohen's unweighted kappa (K) values of ≤ 0 (poor), 0.01-0.20 (slight), 0.21-0.40 (fair), 0.41-0.60 (moderate), 0.61-0.80 (substantial), and 0.81-1 (almost perfect), and weighted kappa (Kw) values of ≤ 0.20 (poor), 0.21-0.40 (fair), 0.41-0.60 (moderate), 0.61-0.80 (good), and 0.81-1.00 (very good) agreement.¹⁵ Multiple logistic regression was used to build prediction models for scores on the mRS and smRSq at Day 90, and to calculate C-indexes. Significant predictors ($P < 0.05$) from the univariate analyses were tested in multiple logistic regression models for their associations with outcomes. The non-significant covariates were removed until all the remaining predictors were statistically significant ($P < 0.05$). Collinearity between variables were checked. Baseline variables included in the models were: age (< 65 vs. ≥ 65 years), sex, estimated prestroke function on mRS (0 vs. 1), baseline NIHSS score, history of atrial fibrillation (AF), diabetes mellitus, hypertension, previous stroke, coronary artery disease, and hypercholesterolemia, use of aspirin/other antiplatelet agent(s), and warfarin/other anticoagulation, and visible early ischemic change and proximal vessel occlusion on imaging.

Correlations between smRSq and mRS at Day 90, and with NIHSS and EQ-5D utility scores at Day 90, were analyzed using Spearman correlation, with the r coefficient graded as 0.2–0.4 (weak), 0.4–0.7 (moderate), and 0.7–1.0 (strong). The treatment effects comparing low-dose alteplase to standard-dose alteplase in the trial were tested using scores derived from smRSq, to compare with the study results generated using mRS. The noninferiority margin was 1.14,^{10,11} that is for the upper boundary of the 95% confidence interval (CI) for the odds ratio (OR) with low-dose alteplase as compared with standard-dose alteplase, of less than 1.14. Single logistic regression was used to test and estimate unadjusted OR of death and disability (mRS 2 to 6). Multiple logistic regression were used for adjusted OR in intention to treat and per protocol populations. For shift analyses of the smRSq scores, ordinal logistic regression was used. The variables adjusted in treatment effect analyses include site, time from symptom onset to randomisation, score as a continuous measure on the NIHSS, age, sex, ethnicity, pre-morbid mRS score (0 or 1), pre-morbid use of aspirin, other antiplatelet agent or warfarin, and any history of stroke, coronary artery disease, diabetes mellitus and atrial fibrillation (AF). Testing was undertaken for the degree of agreement between smRSq and mRS at Day 28 using Kappa (K) and weighted Kappa (Kw), and for the strength of correlations between smRSq or mRS at Day 28, and NIHSS or EQ-5D utility scores at Day 28, using Spearman correlation with the r coefficient (Supplementary Appendix). P values <0.05 were regarded as significant. SAS enterprise 7.1 was used in all analyses.

Data sharing

The authors confirm that the data supporting the findings of this study are available within the article and/or its supplementary materials. Individual participant data used in these analyses can be shared by request from any qualified investigators via the Research Office of The George Institute for Global Health, Australia.

Results

There were 3204 acute ischemic stroke patients with NIHSS scores recorded at baseline, and mRS, smRSq and EQ-5D scores recorded at Day 90. Agreement between smRSq and mRS scores occurred in 2051 (64%) patients ([Supp](#) Table 1, Figure 1), and overall was moderate-good (K 0.57, 95% CI 0.55–0.59, and Kw 0.79, 95% CI 0.78–0.81).

[Supplementary](#) Table 2 shows the variables remained in the prediction models were common to both the smRSq and mRS at Day 90 after successively removing non-significant covariates; these included age (>65 years), AF, diabetes mellitus, pre-morbid symptoms, NIHSS scores and signs of cerebral ischemia on imaging. C-indexes for the model fit were similar for the smRSq and mRS (0.74, 95% CI 0.72-0.76, and 0.75, 95% CI 0.73-0.77, mRS, respectively) (Figure 2).

Concordance was also evident for baseline NIHSS scores (positive correlation; r 0.442, $P < 0.0001$ and r 0.455, $P < 0.0001$, respectively) and EQ-5D utility score (negative correlation; r -0.836, $P < 0.0001$, and r -0.874, $P < 0.0001$, respectively) and smRSq and mRS at Day 90.

Comparisons of the treatment effects using smRSq and mRS are presented in [Supp.](#) Table 3. Both the dichotomous and ordinal outcomes using smRSq were similar to the outcomes from mRS. The unadjusted dichotomous outcome (score of smRSq 2 to 6), which was used to compare with the primary outcome of the alteplase-dose arm of the trial (OR 1.09, 95% CI 0.95-1.25; one sided $P = 0.51$ for noninferiority), occurred in 886 of 1609 patients (55.1%) in the low-dose group and in 863 of 1600 patients (53.9%) in the standard-dose group (OR 1.05, 95% CI 0.91-1.20; one-sided $P = 0.23$ for noninferiority). In the unadjusted shift analysis on smRSq scores comparing low-dose alteplase to standard-dose alteplase, the OR was 0.98 (95% CI 0.87-1.11; $P = 0.02$ for noninferiority) similar to that for mRS shift scores (OR 1.0; 95% CI 0.89-1.13; $P = 0.04$ for noninferiority).

The results for agreement between smRSq and mRS at Day 28, and correlations with NIHSS

and EQ-5D utility score at Day 28, are included in the supplementary appendix.

Discussion

Our study validates the smRSq as a suitable stroke outcome measure by showing comparable scoring to the conventional mRS, similar level of moderate-strong correlations with the NIHSS and EQ-5D, common predictor variables and similar treatment effects when used as trial outcome.

Dennis et al.³ showed similar agreement between the mRS and smRSq using postal or telephone assessment in 225 participants, whilst Yuan et al.⁹ found a higher degree of overall agreement than we have shown in their study of 150 Chinese patients. The factors identified in our predictive models for the smRSq and mRS support other outcome studies.^{16, 17} For example, in a multivariable analysis by Wahlgren et al.,¹⁶ older age, high blood glucose, high NIHSS, and infarction on brain imaging were found to predict poor outcome (mortality or dependency) in patients treated with intravenous alteplase, whilst pre-stroke disability was only associated with mortality. Baseline severity, history of diabetes mellitus, ischemic stroke, and peripheral artery disease have also been reported to predict recovery after disabling ischemic stroke.¹⁷ Katzan et al.¹⁸ showed only a moderate correlation ($r=-0.53$, $p<0.01$) between the mRS and EQ-5D utility score, possibly due to the greater number of patients with mRS scores of 0-2 (75%), which has shown a lower correlation with EQ-5D¹⁹, than in the ENCHANTED¹⁰ (~65%). Another study showed the smRSq had moderate correlation with the physical ($r=0.50$, $P=0.005$) but only slight correlation with the mental components ($r=0.36$, $P=0.048$) of the 12-item short form questionnaire.⁵

More severe strokes (NIHSS scores >10) are associated with higher mRS scores at hospital discharge.²⁰ NIHSS scores at Day 2 are a good predictor of mRS scores >3 at 90 days.²¹ In a study of acute ischemic stroke patients treated with mechanical thrombectomy, NIHSS scores

173 at baseline and hospital discharge were each significantly associated with 90-day mRS scores.²²
174 Another study has shown a similar moderate level of correlation between initial NIHSS and
175 Day 90 smRSq scores ($r = 0.69$, $R^2 = 0.47$, $P < 0.001$)⁵ to our study.

176 The smRSq appears easy to administer and automatically calculates a final score from the
177 structured responses to five questions, whereas the mRS often requires the assessor to make a
178 judgment call in deciding which category best fits a certain grading of disability or level of
179 dependency. While training in the use of the mRS is often used to decrease error, this can be
180 resource intensive for large studies. It is interesting to note that a high percentage of patients
181 who scored 1 or 2 on the mRS scored 3 on smRSq in our study. One explanation could be that
182 a high proportion of ENCHANTED patients experienced acalculia and difficulty managing
183 finances without major motor disability after suffering a left middle cerebral artery stroke. This
184 may have resulted in them answering negatively the first question of the smRSq, resulting in a
185 score ≥ 3 . Another explanation is broader cognitive impairment but we did not collect such
186 information in the study.

187 Our analyses found that similar factors were predictors of smRSq and mRS. This confirms the
188 good correlation between the two scales and re-enforces that they are well-known predictors
189 of poor outcome. Similarly, the correlation between smRSq and mRS is good which is not
190 surprising as both scales correlated similarly with the NIHSS and EQ-5D.

191 In reviewing the treatment effects of the alteplase-dose arm of ENCHANTED, use of the
192 smRSq similarly failed to show that low-dose alteplase was noninferior to standard-dose
193 alteplase with respect to death or disability at Day 90, but was non-inferior with respect to
194 ordinal shift of smRSq scores, which is consistent with those results using mRS.¹⁰ This again
195 reflects good correlation between the two measures, and for the smRSq to provide a comparable
196 assessment of a treatment effect to that on the mRS.

Strengths of this study is the large database of prospectively and systematically assessed patients from a variety of countries and ethnic backgrounds. There are some limitations including that these were post-hoc analyses and that the same outcome assessors rated the mRS and smRSq. However, the Day 90 assessment case report form was structured for sequential recording of the mRS followed by smRSq, and these people were not provided with scoring answers to the smRSq questions. Another issue is that as patients with pre-morbid functional impairment/disability (mRS >1) were excluded from the trial, we are unable to provide an assessment of any influence of this factor on the correlation between the measures. Moreover, the finding of large proportion of patients in the score of 3 using smRSq, similarly shown in the FOCUS trial,²³ suggests distribution of patients across categories may differ between mRS and smRSq, which potentially influenced the results of this study. Finally, as this work pertains to a clinical trial involving acute ischemic stroke patients of predominantly mild-moderate severity, caution may be required in generalizing these results to a more severe patient population or in those with acute intracerebral hemorrhage.

In summary, our study has shown that the smRSq has comparable scoring and construct to the conventional mRS, and provides a useful, reliable and valid outcome measure in the assessment of patients with acute ischemic stroke.

214 **Author contributions**

215 XC undertook analyses and wrote the first draft of the manuscript; CD, JL and CSA interpreted
216 the data; other authors provided critical review; all authors contributed to drafting and take
217 responsibility for the content and integrity of this article.

218 **Funding/Support**

219 The ENCHANTED study received grants from the National Health and Medical Research
220 Council (NHMRC) of Australia (Project Grant numbers 1020462 and 1101113), the Stroke
221 Association of the United Kingdom (TSA 2012/01 and 2015/01), the Ministry of Health and
222 the National Council for Scientific and Technological Development of Brazil (CNPQ:
223 467322/2014-7, 402388/2013-5), the Ministry for Health, Welfare and Family Affairs of the
224 Republic of Korea (HI14C1985). During the completion of this work Maree Hackett was
225 supported by a NHMRC Career Development Fellowship Level 2 (APP1141328) and Craig
226 Anderson a NHMRC Senior Principal Research Fellowship.

227 **Role of the Funders/Sponsors**

228 The funding bodies had no role in the design and conduct of the analyses and interpretation of
229 the data; and preparation, review, or approval of the manuscript

230 **Conflicts of Interest Disclosures**

231 Dr. Anderson reports receiving fees for serving on advisory boards from Amgen, Boehringer
232 Ingelheim, and lecture fees and travel support from Takeda. Dr Chalmers reports research
233 grants and lecture fees from Servier for the ADVANCE trial and post-trial follow-up. Dr.
234 Lavados reports receiving fees for serving on the advisory boards from ANGELS initiative and
235 lectures fees from Boehringer Ingelheim and grant support from Bayer and Boehringer
236 Ingelheim. The other authors report no conflicts of interest.

237 **Acknowledgement**

238 We acknowledge the contribution of the large number of patients, hospital site investigators
239 and coordinators, and central and regional project staff for the ENCHANTED study. We thank
240 EuroQoL Group for providing translations and license for EQ-5D.

For Review Only

Reference

1. Bamford JM, Sandercock PA, Warlow CP, Slattery J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke* 1989;20:828
2. Saver JL, Filip B, Hamilton S, Yanes A, Craig S, Cho M, et al. Improving the reliability of stroke disability grading in clinical trials and clinical practice: the Rankin Focused Assessment (RFA). *Stroke* 2010;41:992-995
3. Dennis M, Mead G, Doubal F, Graham C. Determining the modified Rankin Score after stroke by postal and telephone questionnaires. *Stroke* 2012;43:851-853
4. Wilson JT, Hareendran A, Grant M, Baird T, Schulz UG, Muir KW, et al. Improving the assessment of outcomes in stroke: use of a structured interview to assign grades on the modified Rankin Scale. *Stroke* 2002;33:2243-2246
5. Bruno A, Shah N, Lin C, Close B, Hess DC, Davis K, et al. Improving modified Rankin Scale assessment with a simplified questionnaire. *Stroke* 2010;41:1048-1050
6. Bruno A, Shah N, Akinwuntan AE, Close B, Switzer JA. Stroke size correlates with functional outcome on the simplified modified Rankin Scale questionnaire. *J Stroke Cerebrovasc Dis* 2013;22:781-783
7. Bruno A, Akinwuntan AE, Lin C, Close B, Davis K, Baute V, et al. Simplified modified Rankin Scale questionnaire: reproducibility over the telephone and validation with quality of life. *Stroke* 2011;42:2276-2279
8. Bruno A, Close B, Switzer JA, Hess DC, Gross H, Nichols III FT, et al. Simplified modified Rankin Scale questionnaire correlates with stroke severity. *Stroke* 2013;27:724-727
9. Yuan JL, Bruno A, Li T, Li SJ, Zhang XD, Li HY, et al. Replication and extension of the simplified modified Rankin Scale in 150 chinese stroke patients. *Eur Neurol*

- 2012;67:206-210
10. Anderson CS, Robinson T, Lindley RI, Arima H, Lavados PM, Lee TH, et al. Low-dose versus standard-dose intravenous alteplase in acute ischemic stroke. *N Engl J Med* 2016;374:2313-2323
 11. Huang Y, Sharma VK, Robinson T, Lindley RI, Chen X, Kim JS, et al. Rationale, design, and progress of the Enhanced Control of Hypertension AND thrombolysis stroke study (ENCHANTED) trial: an international multicenter 2 x 2 quasi-factorial randomized controlled trial of low- vs. standard-dose rt-PA and early intensive vs. guideline-recommended blood pressure lowering in patients with acute ischaemic stroke eligible for thrombolysis treatment. *Int J Stroke* 2015;10:778-788
 12. Dolan P, Gudex C, Kind P, Williams A. A social tariff for EuroQol: results from a UK general population survey. 1995
 13. Kind P, Hardman G, Macran S. UK population norms for EQ-5D. 1999
 14. Cohen J. Weighted kappa: Nominal scale agreement with provision for scaled disagreement or partial credit. *Psychol Bull* 1968;70:213-220
 15. Brennan P, Silman A. Statistical methods for assessing observer variability in clinical measures. *BMJ* 1992;304:1491-1494
 16. Wahlgren AN, Ahmed GN, Eriksson RN, Aichner OF, Bluhmki OE, Dávalos OA, et al. Multivariable analysis of outcome predictors and adjustment of main outcome results to baseline data profile in randomized controlled trials: Safe Implementation of Thrombolysis in Stroke-Monitoring Study (SITS-MOST). *Stroke* 2008;39:3316-3322
 17. Hankey GJ, Spiesser J, Hakimi Z, Bego G, Carita P, Gabriel S. Rate, degree, and predictors of recovery from disability following ischemic stroke. *Neurology* 2007;68:1583-1587

- 289 18. Katzan IL, Thompson NR, Lapin B, Uchino K. Added value of patient-reported
290 outcome measures in stroke clinical practice. *J Am Heart Assoc* 2017;6
- 291 19. Rangaraju S, Haussen D, Nogueira RG, Nahab F, Frankel M. Comparison of 3-month
292 stroke disability and quality of life across modified Rankin Scale categories. *Interv*
293 *Neurol* 2017;6:36-41
- 294 20. Mihindu E, Mohammed A, Smith T, Brinster C, Sternbergh WC, 3rd, Bazan HA.
295 Patients with moderate to severe strokes (NIHSS score >10) undergoing urgent carotid
296 interventions within 48 hours have worse functional outcomes. *J Vasc Surg*
297 2019 ;69 :1471-1481
- 298 21. De Raedt S, Brouns R, De Smedt A, Aries MJ, Uyttenboogaart M, Luijckx GJ, et al.
299 The sNIHSS-4 predicts outcome in right and left anterior circulation strokes. *Clin*
300 *Neurol Neurosurg* 2013;115:729-731
- 301 22. Costalat V, Lobotesis K, Machi P, Mourand I, Maldonado I, Heroum C, et al.
302 Prognostic factors related to clinical outcome following thrombectomy in ischemic
303 stroke (RECOAST study): 50 patients prospective study. *Eur J Radiol* 2012;81:4075-
304 4082
- 305 23. Dennis M, Mead G, Forbes J, Graham C, Hackett M, Hankey GJ, et al. Effects of
306 fluoxetine on functional outcomes after acute stroke (FOCUS): A pragmatic, double-
307 blind, randomised, controlled trial. *The Lancet*. 2019;393:265-274

308

Table 1. Correlation between smRSq and mRS scores at Day 90

mRS	smRSq							
	0	1	2	3	4	5	6	Total
0	704 (88.0)	42 (5.3)	11 (1.3)	42 (5.3)	1 (0.1)	-	-	800
1	311 (42.4)	266 (36.3)	56 (7.6)	97 (13.2)	3 (0.4)	-	-	733
2	38 (8.0)	75 (15.8)	167 (35.2)	174 (36.6)	16 (3.4)	5 (1.1)	-	475
3	6 (1.5)	8 (2.0)	22 (5.6)	285 (72.7)	52 (13.3)	19 (4.6)	-	392
4	5 (1.6)	2 (0.6)	2 (0.6)	36 (11.3)	162 (50.9)	111 (34.9)	-	318
5	2 (1.1)	-	-	2 (1.1)	15 (8.5)	157 (89.2)	-	176
6	-	-	-	-	-	-	310 (100)	310
Total	1066	393	258	636	249	292	310	3204*

mRS denotes modified Rankin Scale, smRSq simplified modified Rankin Scale questionnaire.

Kappa statistic 0.57 (95% confidence interval [CI] 0.55–0.59) and weighted Kappa statistic 0.79 (95% CI 0.78–0.81)

*3310 patients were randomized into the alteplase dose arm, of which 13 were excluded; another 93 patients were excluded from these analyses due to missing mRS or smRSq data.

Table 2. Independent predictors of smRSq and mRS at Day 90

Variable	smRSq (C=0.740, 95% CI 0.723-0.757)			mRS (C=0.751, 95% CI 0.734-0.767)		
	OR	95% CI	p-Value	OR	95% CI	p-Value
Age >65	1.47	1.26-1.71	<0.0001	1.33	1.13-1.56	0.0005
Atrial fibrillation	1.43	1.16-1.77	0.0009	1.29	1.04-1.59	0.019
Diabetes mellitus	1.25	1.03-1.51	0.0245	1.37	1.13-1.66	0.002
Pre-stroke grade of physical function*	2.21	1.79-2.72	<0.0001	2.24	1.82-2.77	<0.0001
NIHSS	1.14	1.12-1.16	<0.0001	1.16	1.14-1.17	<0.0001
Signs of cerebral ischemia on imaging	1.56	1.30-1.88	<0.0001	1.42	1.18-1.71	0.0002

C denotes Concordance Index, CI confidence interval, mRS modified Rankin Scale, NIHSS National Institutes of Health Stroke Scale, OR odds ratio, smRSq simplified modified Rankin Scale questionnaire

*pre-morbid estimated level of physical function with symptoms, based on a score of 1 on the mRS; the comparison was 1 vs. 0

Significant predictors (P<0.05) from the univariate analyses which were tested in multiple logistic regression models were: sex, history of hypertension, previous stroke, coronary artery disease, hypercholesterolemia, use of aspirin/other antiplatelet agent(s), use of warfarin/other anticoagulation and proximal vessel occlusion. Significance level of stay in the models was P<0.05.

Table 3. Comparison of treatment effects using mRS and smRSq in the alteplase-dose arm of the ENCHANTED trial

Outcome	smRSq			mRS		
	OR	95% CI	P-value*	OR	95% CI	P-value*
Death or disability: scores 2 to 6 [‡]						
—Unadjusted	1.05	0.91–1.20	0.23	1.09	0.95–1.25	0.51
—Adjusted [‡]	1.06	0.91–1.23	0.34	1.13	0.97–1.31	0.88
—Adjusted in per protocol population [§]	1.05	0.89–1.23	0.30	1.13	0.96–1.32	0.89
Shift analyses of scores 0 to 6 [‡]						
—Unadjusted	0.98	0.87–1.11	0.02	1.00	0.89–1.13	0.04
—Adjusted [‡]	0.97	0.85–1.10	0.01	0.99	0.88–1.13	0.03
—Adjusted in per protocol population [§]	0.95	0.84–1.09	0.01	1.00	0.88–1.14	0.05

CI denotes confidence interval, mRS modified Rankin Scale, OR odds ratio, smRSq simplified modified Rankin Scale questionnaire

*Noninferiority margin was 1.14 (i.e. an upper boundary of the 95% CI for the OR with low-dose alteplase as compared with standard-dose alteplase of less than 1.14).

[‡]ORs were estimated from logistic regression models. Each OR indicates the odds of death or disability (mRS 2 to 6). An OR greater than 1 favors standard-dose alteplase. Adjustment for site, time from stroke onset to randomisation, score as a continuous measure on the National Institutes of Health stroke scale (NIHSS), age, sex, ethnicity, pre-morbid score of 0 or 1 on the mRS, pre-morbid use of aspirin, other antiplatelet agent or warfarin, and any history of stroke, coronary artery disease, diabetes mellitus and atrial fibrillation.

[‡]ORs were estimated from ordinal logistic regression models. Each OR indicates the odds of an increase of 1 in the mRS score. An OR greater than 1 favors standard-dose alteplase. Adjustment for same variables as in logistic regression models above.

[§]Per protocol population excluded patients who have one or more of the following protocol violations: age <18 years; final diagnosis not acute ischemic stroke; final diagnosis unknown/uncertain because of missing source documents or neuroimaging; baseline systolic blood pressure >185 mmHg; randomized >4.5 hours; failure to receive alteplase at either the correct bolus or infusion dose; failure to obtain a blind assessment of the 90-day outcome.

Figures legend

Figure 1. Bubble plot of agreement between smRSq and mRS at Day 90. Area of bubbles represent the count at each score.

Figure 2. ROC curves for predictive models of mRS and smRSq at Day 90.

For Review Only


Figure 1. Agreement between smRSq and mRS at Day 90

Figure 2. ROC curves for the predictive models of mRS and smRSq at Day 90

